

1  
2  
3  
4  
5

# **Heliospheric Magnetic Field 1835-2009**

Leif Svalgaard<sup>1</sup> and Edward W. Cliver<sup>2</sup>

<sup>1</sup>Stanford University, HEPL, Cedar Hall, Via Ortega, Stanford, CA 94305-4085

<sup>2</sup>Space Vehicles Directorate, Air Force Research Laboratory, Hanscom AFB, MA

01731-3010

6 **Abstract.** We use recently acquired geomagnetic archival data to extend our long-term  
7 reconstruction of the HMF strength. The 1835-2009 HMF series is based on an updated  
8 and substantiated *IDV* series from 1872-onwards and on Bartels' extension, by proxy, of  
9 his *u*-series from 1835-1871. The new *IDV* series, termed IDV09, has excellent  
10 agreement ( $R^2 = 0.98$ ; RMS = 0.3 nT) with the earlier IDV05 series, and also with the  
11 negative component of Love's extended (to 1905)  $D_{st}$  series ( $R^2 = 0.91$ ). Of greatest  
12 importance to the community, in an area of research that has been contentious,  
13 comparison of the extended HMF series with other recent reconstructions of solar wind *B*  
14 for the last ~100 years yields a strong consensus between series based on geomagnetic  
15 data. Differences exist from ~1900-1910 but they are far smaller than the previous  
16 disagreement for this key interval of low solar wind *B* values which closely resembles  
17 current solar activity. Equally encouraging, a discrepancy with an HMF reconstruction  
18 based on  $^{10}\text{Be}$  data for the first half of the 20th century has largely been removed by a  
19 revised  $^{10}\text{Be}$ -based reconstruction published after we submitted this paper, although a  
20 remaining discrepancy for the years ~1885-1905 will need to be resolved.

## 21 **1. Introduction**

22 In *Svalgaard and Cliver* [2005] we introduced the InterDiurnal Variability (*IDV*) index  
23 for a given geomagnetic observatory ('station') as the average difference without regard  
24 to sign, from one day to the next, between hourly mean values of the Horizontal  
25 Component,  $H$ , and measured one hour after midnight. The average should be taken over  
26 a suitably long interval of time, such as one year, to eliminate various seasonal  
27 complications.

28 *IDV* has the useful property of being independent of solar wind speed and is highly  
29 correlated with the near-Earth Heliospheric Magnetic Field (HMF) strength  $B$ . Thus once  
30 *IDV* is determined, solar wind  $B$  is known as well. *Svalgaard and Cliver* [2005] used *IDV*  
31 augmented with *Bartels'* [1932]  $u$ -measure to reconstruct the HMF strength for the years  
32 1872-2004.

33 Here we report on an extension of the *IDV* index for a longer time interval (1835-2009),  
34 using many more stations. The inclusion of more data is particularly important for the  
35 years from 1890-1909 for which the initial version of the index (*IDV05*) was based on  
36 observations from only one station before 1901 and four more stations from 1903. An  
37 important aspect of *IDV09* is that it includes recent years with index values at the same  
38 level as the very low values in 1901-1902, thus allowing the correlation between *IDV* and  
39 the magnitude of the near Earth HMF to be extended to such low values without  
40 extrapolation. With this correlation, we infer HMF  $B$  for years prior to the space age and  
41 compare our  $B$  values with those obtained by other investigators using geomagnetic or  
42 cosmic ray data.

## 43 2. Analysis

### 44 2.1 Derivation of IDV09

45 Our determination of IDV09 is essentially identical to that of IDV05 except for the  
46 inclusion of more data. In *Svalgaard and Cliver* [2005] we emphasized that *IDV* is a  
47 modern version of the *u*-measure building on ideas of a century ago [*Moos*, 1910]. *Kertz*  
48 [1958], *Mayaud* [1980], and *Svalgaard* [2005] suggested using only night-time values to  
49 avoid contamination by the regular diurnal variation,  $S_R$ . We followed their lead but  
50 further limited the time interval to only one hour following local midnight and  
51 constructed the interdiurnal variability index (*IDV*) for a given station as the unsigned  
52 difference between two consecutive days of the average value over the interval of a *H*  
53 component measured in nT. The individual unsigned differences were then averaged over  
54 longer intervals, *e.g.*, one full year (minimizing various geometric and seasonal effects,  
55 *e.g.* the semiannual non-solar variation due to the tilt of the Earth's dipole – a plot of 27-  
56 day Bartels Rotation values of *IDV* can be found in *Svalgaard* [2009]).

57 Since 2005, we have been collecting, digitizing, quality controlling, and correcting  
58 (where needed) hourly historical geomagnetic data from individual observatories as well  
59 as from World Data Centers [there is, as yet, no mechanism available to individual  
60 researchers for injecting new or corrected data into the World Data Centers or into  
61 various National Depositories, so we offer the data to interested researchers upon  
62 request]. Here we use these newly-acquired data to substantiate the *IDV*-index, which is  
63 especially important for the first ~30 years of the time series (1872-1902), during which  
64 IDV05 was based solely on *Schmidt's* [1926] and *Bartels'* [1932] *u*-measure from 1872-  
65 1889, on Potsdam observations from 1890-1902, plus Cheltenham for 1901-1902, and

66 Honolulu for 1902. In contrast, IDV09 is based on four times as many “station years”  
67 (143 vs. 34) for this 31-yr interval as detailed in Table E1 of the Electronic Supplement.  
68 We update the time series by adding the index values for 2004-2009. These latter years  
69 are significant because the yearly-averages of  $B$  observed in 2007-2009 are the lowest  
70 observed during the space age. They lie at the lower endpoint of the correlation between  
71 yearly averages of observed  $B$  and  $IDV$ .

72 Table 1 contains a list of the 71 stations (including replacement stations) used to compute  
73 IDV09 (versus 14 for IDV05). A comprehensive list of the data coverage and the data  
74 values for the individual stations used in this study is given in Table E1 in the Electronic  
75 Supplement. All raw data is available from the authors (LS) upon request.

#### 76 **2.1.1. Latitude Normalization**

77 For IDV05, we normalized  $IDV$  values for a given station with Corrected Geomagnetic  
78 Latitude,  $M$ , to those of Niemegek (NGK) [as Bartels did for the  $u$ -measure] using

$$79 \quad IDV_{\text{norm}} = IDV_{\text{raw}} / (1.324 \cos^{0.7}(M)) \quad (1)$$

80 Here we have retained this relationship for stations with  $|M| < 51^\circ$  because it still fits the  
81 data for the additional stations. At significantly higher latitudes, the index becomes  
82 strongly contaminated by auroral zone activity [see Figure 2 of *Svalgaard and Cliver*,  
83 2005, and we recommended not using such stations, *e.g.*, the long-running station  
84 Sodankylä, SOD (used by *Lockwood et al.* [2009]). For IDV09, we relax this restriction  
85 slightly [by a few degrees for a few stations, indicated in Table 1 using a constant,  
86 empirical normalization divisor of 1.1 for these stations, instead of the divisor in equation  
87 (1)]. A value larger than  $\sim 0.95$  for  $|M| \geq 51^\circ$  indicates some contamination by auroral

88 zone activity. We have not attempted to further quantify the latitude dependence of the  
89 contamination, but simply use an average value for the few stations slightly above 51°.  
90 We do this to accommodate changes in  $M$  with time which for some stations can exceed  
91 several degrees<sup>1</sup> and to include a few long-running stations just above 51°. Figure 1  
92 shows the adopted normalization divisor as a function of  $M$  for the 71 stations used in the  
93 present study. Different symbols denote the divisor values for the years 1800, 1900, and  
94 2000, showing the sensitivity of  $IDV$  to changes in latitude. The normalization divisor  
95 was calculated for the centroid of the latitudes for the actual data coverage for each  
96 station. If we did not normalize, the presence of data gaps [of which there are many]  
97 would produce discontinuities in the composite series.

### 98 **2.1.2. Effect of Hourly Means versus Hourly Values on $IDV$**

99 Early magnetometer data were taken [and/or reported] as readings once an hour rather  
100 than as the hourly mean that Adolf Schmidt advocated in 1905 and that was widely and  
101 rapidly adopted. In *Svalgaard & Cliver* [2005] we showed that although the variance of  
102 single values is larger than for averages, the overall effect on  $IDV$  was small (at most a  
103 few percent)<sup>2</sup>. The two long-running series POT-SED-NGK and PSM-VLJ-CLF afford a  
104 convenient additional test of this: POT changed from values to means with the 1905  
105 yearbook, but CLF changed much later, with the 1972 yearbook, so we can directly

---

<sup>1</sup> We expect only a very *weak* influence in the basic response of the Ring Current [see section 2.1.5] to the change of the Earth's magnetic dipole moment [as per *Glassmeier et al.*, 2004] over the interval in question, and so have not attempted to correct for this.

<sup>2</sup> This effect is significant for the  $IHV$  index, but for that case, correction of the effect is straightforward [*Svalgaard and Cliver*, 2007b].

106 compare the (raw – uncorrected in any way) *IDV*-values for the two series (Figure 2). It is  
107 evident that the change from hourly instantaneous values to hourly means did not  
108 introduce any sudden changes in *IDV* at the times of the transitions. The Japanese station  
109 at KAK changed from values to means in 1955. The ratio between raw *IDV* for KAK and  
110 SED-NGK (crosses on Figure 2) also does not show any change in 1955. The American  
111 stations CLH and HON changed to means with the 1915 yearbook. Comparison [Figure  
112 3] over a 24-year interval centered on 1915 with the stations VLJ and DBN, which did  
113 not change sampling procedure, also shows no detectable change in *IDV* due to the  
114 change in sampling: the ratio between average CLH-HON and VLJ-DBN is 1.0792  
115 before 1915 and 1.0792 after the change to hourly means in 1915. We conclude that  
116 changes are too small to justify attempting *ad-hoc* correction based on extrapolation of  
117 modern data.

### 118 2.1.3. Using the *u*-measure before 1872

119 Julius *Bartels* [1932] compiled the *u*-measure from the interdiurnal variability of the  
120 Horizontal Component, *H*, from hourly or daily values from several observatories  
121 operating from 1872 onwards as described in his paper. He wrote, “Before 1872, no  
122 satisfactory data for the calculation of interdiurnal variabilities are available”, but “more  
123 for illustration than for actual use”, he attempted to extend the series backwards to 1835.  
124 For this he used the “Einheitliche Deklinations-Variationen”<sup>3</sup>, *E*, of *Wolf* [1884] and the  
125 “summed ranges”, *s*, derived from the mean diurnal variation of *H* at Colaba (Bombay)  
126 due to *Moos* [1910]. He derived regression formulae relating *E* and *s* to *u* for times after

---

<sup>3</sup> Unified Declination Variations

127 1872 and used them to synthesize values of  $u$  for the earlier years<sup>4</sup>; giving  $s$  double the  
128 weight of  $E$ . Bartels justified this by showing that for the annual means 1872-1901, the  
129 values of  $u$  derived from  $H$  and the values of  $s$  have a high linear correlation coefficient.  
130 We have extended his analysis by calculating the correlation between  $IDV$  and the  
131 Summed Ranges for 1872-1905 finding a correlation coefficient of 0.86. Figure 4 shows  
132 the agreement between observed  $IDV$  [red] and that calculated from  $s$  [blue].

133 Furthermore, as shown in *Svalgaard and Cliver* [2005] there is a good linear correlation  
134 between  $IDV$  [or HMF  $B$  derived from it] and the square root of the sunspot number,  $R$ .  
135 The main sources of the equatorial components of the Sun's large-scale magnetic field  
136 are large active regions. If these active regions emerge at random longitudes, their net  
137 equatorial dipole moment will scale as the square root of their number. Thus their  
138 contribution to the average HMF strength will tend to increase as  $R^{1/2}$  (for a detailed  
139 discussion, see *Wang and Sheeley* [2003] and *Wang et al.* [2005]).

140 To the extent that the  $u$ -measure before 1872 can be taken as a geomagnetic-based  
141 measure of the sunspot number, it is therefore to be expected that the  $u$ -measure also will  
142 serve as a proxy for  $IDV$ . This estimate will be independent of any assumptions about the  
143 constancy of the calibration of the sunspot number (*c.f.* the difference between the Zürich  
144 Sunspot Number and the Group Sunspot Number [*Hoyt et al.*, 1994]).

145 Figure 5 shows that  $IDV$  can also be directly inferred from the daily range,  $rY$ , of the East  
146 component [equivalent to the Declination for this purpose] of the geomagnetic field and  
147 that therefore, again, that the  $u$ -index before 1872 [strongly influenced by the range of the

---

<sup>4</sup> From  $E$  and  $s$ , we calculate a value of 0.72 for the value for  $u$  for 1857.5 using the formulae given by Bartels.



148 daily variation] can be used for estimation of *IDV*, albeit with slightly less accuracy than  
149 after 1872. This conclusion may seem at variance [and did surprise us] with our initial  
150 decision to use only night-time data in the derivation of *IDV*, but emerges naturally [and  
151 inescapably] after our analysis had shown that *IDV* derived without any dependence on  
152 daytime data is comparable to *IDV* derived from daily ranges because of the strong  
153 dependence of both on the sunspot number. This is clearly demonstrated in Figure 6 that  
154 shows raw *IDV* calculated for PSM-VLJ-CLF and POT-SED-NGK determined from  
155 night-time differences (blue) and daytime differences (red). This realization opens the  
156 door for use of 19<sup>th</sup> geomagnetic stations that only observed during the day as long as the  
157 observations were made at fixed hours.

158 For the reasons given above, we find that *IDV* can be estimated with confidence from  
159 Bartels' *u*-measure also before 1872, justifying our reconstruction of HMF *B* since 1835,  
160 with expected refinement of the reconstruction upon further digitization of 19<sup>th</sup> century  
161 geomagnetic yearbooks.

#### 162 **2.1.4. The *IDV*-index 1835-2009**

163 From the ~1,375,000 daily differences [3775 station-years] derived from the stations in  
164 Table 1 we construct the *IDV*-index shown in Figure 7, with individual station curves in  
165 grey. The composite (red) curve is the mean of the median and average values for each  
166 year, while before 1872 the dashed curve shows *IDV* estimated from *u*. Also shown (blue  
167 curve) is the number of stations contributing to the mean. The large number of stations  
168 from 1957 on does not add further significance to the composite, but only serves to  
169 establish the range of scatter of the values.

170 It is evident that *IDV* from only a single station (provided that not too much data is  
171 missing either because the recording went off-scale or as a result of other problems) does  
172 not differ much from the mean of many stations; the standard deviation of *IDV*-values for  
173 all stations for a given year is less than 1 nT or about 9%. This means that only a few  
174 [good] stations are needed for a robust determination of *IDV*. This conclusion, of course,  
175 only emerges after the spread of *IDV*-values has first been shown to be small. The  
176 standard error of the mean of more than fifty stations is 0.1 nT.

177 Figure 8 shows that the differences between *IDV05* and *IDV09* are slight, and due to the  
178 additional data since 1880. During the period of overlap (1872-2003, 2004 was only  
179 partial), the two time series agree within an RMS of 0.33 nT or 3%. The coefficient of  
180 determination for the correlation between *IDV09* and *IDV05* is  $R^2 = 0.984$ . *IDV* is a  
181 robust index.

#### 182 **2.1.5. Physical Interpretation of *IDV*: Measure of the Energy in the Ring Current**

183 In *Svalgaard and Cliver* [2005] we reported that *IDV* is closely correlated with the  
184 negative part of the  $D_{st}$ -index based on data back to 1932 [*Karinen and Mursula*, 2005].  
185 In *Svalgaard and Cliver* [2006] we extended that relationship back to 1905 using the 100-  
186 year  $D_{st}$ -series derived by *J. Love* [2006, 2007], and confirm it here using *IDV09*. Yearly  
187 averages of  $D_{st}$  [scaled to Kyoto  $D_{st}$ ; we use  $D_{st}$  here in a generic sense without  
188 distinguishing between different derivations of the underlying physical measure sought  
189 captured by  $D_{st}$ ] when the hourly value was negative were computed and found to be  
190 strongly correlated with *IDV* [ $R^2 = 0.91$ ]:  $IDV = -0.45 (D_{st} < 0)$ . Figure 9 compares *IDV09*  
191 and *IDV* computed from  $D_{st}$ . The good match suggests that *IDV* is a measure of the same  
192 physical reality as negative  $D_{st}$ , namely the energy in the Ring Current, which then in turn

193 seems to be controlled by HMF  $B$ : ( $D_{st}<0$ ) =  $4.81 B - 9.41$  [ $R^2 = 0.84$ ], and we can then  
 194 also use  $D_{st}$  to determine the HMF strength:  $B = 2.70 - 0.1736 (D_{st}<0)$ . Schmidt [1926]  
 195 actually suggested that in the definition of the  $u$ -measure it would be slightly better to  
 196 only use the negative differences between consecutive days.

## 197 **2.2. Using IDV09 to Calculate HMF Strength, 1835-2009**

198 Since the 2005 definition paper, lower values of HMF strength,  $B$ , have improved the  
 199 dynamic range (and thus the statistical significance) of the correlation between  $IDV$  and  
 200  $B$ . An approximate linear correlation was found, but there is no *a priori* reason the  
 201 relationship would be strictly linear. In addition, it has been argued [Lockwood *et al.*  
 202 2006] that  $B$  should be taken as the independent variable instead of  $IDV$ . We showed in  
 203 Svalgaard and Cliver [2006] that it does not make much difference which way the  
 204 correlation is evaluated. In the end, the RMS difference [0.4 nT or less than ~10%]  
 205 between HMF  $B$  observed *in situ* near the Earth<sup>5</sup> and inferred from  $IDV$  is what matters.  
 206 The average coefficients for the linear correlation performed four ways (average, median,  
 207 and for each: direct and inverse) are

$$208 \quad B \text{ (nT)} = (2.06 \pm 0.21) + (0.441 \pm 0.021) IDV \quad (R^2 = 0.869) \quad (2)$$

209 The equivalent power law dependence comes to

$$210 \quad B \text{ (nT)} = (1.33 \pm 0.07) IDV^{0.689 \pm 0.023} \quad (R^2 = 0.905) \quad (3)$$

211 The adopted values for  $B$  inferred from IDV09 given in Table 2 are the mean values  
 212 calculated using these two relationships. Table E3 in the Electronic Supplement  
 213 summarizes the coefficients for all correlations. The ‘error bars’ quoted are not a measure

---

<sup>5</sup> Using hourly averages from the OMNI dataset for historical data and from ACE for near real-time recent data.

214 of the statistical significance of the correlations in a strict sense, but are solely indicative  
215 of the range or variability of the various coefficients.

216 Figure 10 shows the values for HMF  $B$  inferred from  $IDV$  from 1835 to the present (blue  
217 curve) and  $B$  measured by spacecraft (red curve). A 4<sup>th</sup>-order polynomial fit suggests a  
218 ~100 year Gleissberg cycle. Cycle 23 looks remarkably like cycle 13, including the very  
219 deep solar minimum following both cycles, likely presaging a weak cycle 24 as predicted  
220 from the solar polar fields [Svalgaard *et al.*, 2005; Schatten, 2005]. It is clear that we are  
221 returning to conditions prevailing a century ago. It seems likely that other solar  
222 parameters such as Total Solar Irradiance [Fröhlich, 2009; Steinhilber *et al.*, 2009] and  
223 cosmic ray modulation [Steinhilber *et al.*, 2010] are reverting to similar conditions with  
224 possible implications for the climate-change debate.

## 225 **2.3. Comparison of IDV09-based $B$ with Other Recent Reconstructions**

### 226 **2.3.1. Consilience of Reconstructions Based on Geomagnetic Data.**

227 Reconstructions of HMF  $B$  have been discordant in the past [e.g. Svalgaard, 1977,  
228 referenced in Schatten *et al.*, 1978; Andreasen, 1997; Lockwood *et al.*, 1999, 2006;  
229 Svalgaard and Cliver, 2005, 2006, 2007b]. The realization [Svalgaard *et al.*, 2003] that  
230 geomagnetic indices can be constructed that have different dependencies on  $B$  and solar  
231 wind speed ( $V$ ) has enabled robust determinations of both  $V$  [Svalgaard and Cliver,  
232 2007b; Rouillard *et al.*, 2007; Lockwood *et al.*, 2009] and  $B$  [Svalgaard and Cliver, 2005,  
233 2006; Lockwood *et al.*, 2009] that have converged to a common, well-constrained dataset.  
234 Progress has been swift and Figure 11 shows the convergence of HMF  $B$  determined by  
235 Lockwood *et al.* [2009] to the values determined from  $IDV$  [Svalgaard and Cliver, 2005];

236 this paper]. The *Lockwood et al.* [2009, and references therein] reconstruction is still  
237 discordant from ours for a few years during solar cycle 14, but apart from that, the  
238 agreement is quite remarkable and the issues seem resolved.

239 Figure 12 details the evolution of the various determinations of  $B$  since the influential,  
240 but now superseded, *Lockwood et al.* [1999] paper. It is clear that we now possess the  
241 methodology to infer  $B$  with good accuracy as far back as continuous geomagnetic  
242 records of  $H$  reach. A concerted effort of digitization of 19<sup>th</sup> century yearbook records  
243 would promise to further improve our knowledge of the magnetic field in the heliosphere.  
244 *Svalgaard and Cliver* [2007a] argued for a floor in yearly averages of solar wind  $B$  which  
245 was approached at every 11-yr minimum and represented the ground-state of the Sun  
246 during extended minima such as the Maunder Minimum. With the larger dynamic range  
247 afforded by the current minimum, we can now refine the value of the floor to be closer to  
248 the  $\sim 4$  nT observed during 2008 and 2009 [see also *Owens et al.*, 2008], returning to the  
249 values inferred for 11-yr minima during the previous Gleissberg minimum at the turn of  
250 the 20<sup>th</sup> century.

### 251 2.3.2. Comparison with <sup>10</sup>Be-based Reconstructions

252 *McCracken* [2007] spliced together <sup>10</sup>Be data, ionization-chamber cosmic ray data  
253 (calibrated with balloon flight data), and neutron monitor cosmic ray data to produce an  
254 ‘equivalent’ neutron monitor count series covering the entire interval 1428-2005, and  
255 inverted the series for  $B$  in order to express the data in terms of the HMF  $B$ . In Figure 13  
256 we compare his series for HMF  $B$  with the ‘consensus’  $B$  from geomagnetic data.

257 In McCracken's time series for  $B$ , a large step-like change (1.7 nT; from 3.5 nT to 5.2 nT;  
258 the largest jump in the entire ~600-year record) occurs between the 1944 and 1954  
259 sunspot minima flanking cycle 18. No such corresponding change is observed in the  
260 concordant reconstructions of *Svalgaard and Cliver* [2005; this paper], *Rouillard et al.*  
261 [2007] and *Lockwood et al.* [2009], nor in  $B$  calculated from the quantity  $BV$  deduced by  
262 *Le Sager and Svalgaard* [2004] using either  $V$  of *Svalgaard and Cliver* [2006] or of  
263 *Rouillard et al.* [2007], or in  $B$  deduced from  $D_{st}$ .

264 *Muscheler et al.* [2007] discuss the uncertainties with the balloon-borne data that form  
265 the basis for McCracken's calibration of the composite equivalent neutron monitor data  
266 before 1951. The strong geomagnetic evidence argues that **this particular** calibration of  
267 the pre-neutron monitor cosmic ray reconstruction is not on a firm footing. We suggest  
268 that part of the reason for the disagreement might lie with the calibration and splicing  
269 together of the disparate cosmic ray datasets.

270 After our paper was submitted, we were pleased to read a paper by *Steinhilber et al.*  
271 [2010] in which a new  $^{10}\text{Be}$ -based reconstruction has moved closer to our reconstruction,  
272 to that of *Rouillard et al.* [2007], and to that of *Caballero-Lopez et al.* [2004] with  
273 diffusion coefficient depending inversely of  $B^2$  ( $a \sim 2$ ). The reconstruction of *Steinhilber*  
274 *et al.* [2010] still differs somewhat with the geomagnetic based reconstructions,  
275 especially for the ~1880-1900 interval [Figure 14] and, just like the previous discrepancy,  
276 this will need to be resolved. We suggest that if the sharp dip around ~1895 is not borne  
277 out by further investigation, the magnitude of earlier excursions to very low values may  
278 also be in doubt. Figure 15 shows  $IDV$  for all stations for the interval 1880-1920 and does

279 not support the marked decrease around ~1895. It is unlikely that data from further  
280 stations will change that conclusion.

### 281 3. Summary and Discussion

282 We have extended our 1872-2004 HMF time series [Svalgaard and Cliver, 2005] to the  
283 years 1835-2009 [Figure 10]. The 1835-1871 interval is based on Bartels'  $u$ -measure,  
284 which he extended from 1871 back to 1835 using Wolf's [1884] Declination index based  
285 on several European stations and Moos' [1910] Summed Ranges from Colaba. The 1872-  
286 2009 interval is based on the  $IDV$ -index, with significantly more data for the early years  
287 (1872-1910). The forward extension of the HMF series through 2009 is important  
288 because the years 2007-2009 witnessed the lowest annual averages of  $IDV$  during the  
289 space age. For the time of overlap between the re-evaluated  $IDV$ -index ( $IDV09$ ) and  
290  $IDV05$ , the difference is very small, testifying to the robustness of the index.

291 A comparison of  $IDV09$ -based HMF strength with those obtained by other investigators  
292 using various combinations and permutations of geomagnetic indices revealed a pleasing  
293 agreement [Figure 11] in what had been previously a contentious field of research  
294 [Figures 12 and 13]. The technique proposed by Svalgaard *et al.* [2003] and adopted by  
295 Rouillard *et al.* [2007] to use indices with different dependencies on  $B$  and  $V$  to separate  
296 these variables has proven out and allowed the vast storehouse of hourly and daily data to  
297 be brought to bear. In particular, the  $B$  values deduced and cross-checked [Le Sager and  
298 Svalgaard, 2004] by this method have substantiated the approach made possible by the  
299  $IDV$ -index as well as, as we suggested in Svalgaard and Cliver [2005] and have  
300 confirmed here, the negative component of the  $D_{st}$ -index [Figure 9]. We conclude that the  
301 long-term variation of heliospheric  $B$  is firmly constrained during the time for which it is

302 based on hourly values of  $H$ , and that current values at the solar minimum between cycles  
303 23 and 24 are back to where they were 108 years ago at the solar minimum between  
304 cycles 13 and 14.

305 Although the recent reconstruction of  $B$  based on  $^{10}\text{Be}$  data [Steinhilber *et al.*, 2010]  
306 generally agrees well with the geomagnetic-based reconstruction there is disagreement  
307 for the decade just prior to 1900 [Figure 14]. Further examination of these years is critical  
308 because they present the only example during the 175 year interval of geomagnetic-based  
309  $B$  where the floor [Svalgaard and Cliver, 2007a] in the solar wind is challenged by  
310 Steinhilber *et al* [2010], a challenge not supported by the geomagnetic evidence [Figure  
311 15].



312 **Acknowledgements**

313 Geomagnetic data has been downloaded from the World Data Centers for Geomagnetism  
314 in Kyoto, Japan, and Copenhagen, Denmark [now defunct], and from INTERMAGNET  
315 at [http://www.intermagnet.org/Data\\_e.html](http://www.intermagnet.org/Data_e.html). The research results presented in this paper  
316 rely on the data collected at magnetic observatories worldwide, and we thank the national  
317 institutions that support them. We also recognize the role of the INTERMAGNET  
318 program in promoting high standards of magnetic observatory practice. We thank the  
319 many people worldwide who have helped us with collection of data and metadata in  
320 addition to what is available from public sources. We thank Vladimir Papitashvili for the  
321 program to calculate corrected geomagnetic coordinates using the GUFM1 coefficients  
322 (courtesy of Catherine Constable). The OMNI dataset was downloaded from  
323 <http://omniweb.gsfc.nasa.gov/>. Real-time ACE interplanetary data is downloaded from  
324 <http://www.swpc.noaa.gov/ftpmenu/lists/ace2.html>.

325 **References**

326 **Andreasen, G. K. (1997), Reconstruction of past solar wind variations: Inversion of the**  
327 **geomagnetic response at Godhavn, *J. Geophys. Res.*, *102(A4)*, 7025,**  
328 **doi:10.1029/96JA03161.**

329 Bartels, J. (1932), Terrestrial-magnetic activity and its relations to solar phenomena, *Terr.*  
330 *Magn. Atmos. Electr.*, *37*, 1.

331 Caballero-Lopez, R. A., H. Moraal, K. G. McCracken, and F. B. McDonald (2004), The  
332 heliospheric magnetic field from 850 to 2000 AD inferred from <sup>10</sup>Be records, *J. Geophys.*  
333 *Res.*, *109*, A12102, doi:10.1029/2004JA010633.

334 Fröhlich, C. (2009), Evidence of a long-term trend in total solar irradiance, *Astron. &*  
335 *Astrophys.*, *501(3)*, L27-L30, doi:10.1051/0004-6361/200912318.

336 Glassmeier, K., J. Vogt, A. Stadelmann, and S. Buchert (2004), Concerning long-term  
337 geomagnetic variations and space climatology, *Ann. Geophys.*, *22(10)*, 3669, Sref-  
338 ID:1432-0576/ag/2004-22-3669.

339 **Hoyt, D. V., K. H. Schatten, and Elizabeth Nesme-Ribes (1994), The one hundredth year**  
340 **of Rudolf Wolf's death: Do we have the correct reconstruction of solar activity?**  
341 ***Geophys. Res. Lettrs.*, *21 (18)*, 2067, doi:10.1029/94GL01698.**

342 Joos, G., J. Bartels, and P. Ten Bruggencate (1952), Landolt-Börnstein: Zahlenwerte und  
343 Funktionen aus Physik, Chemie, Astronomie, Geophysik und Technik, in *Astronomie und*  
344 *Geophysik*, vol. XVIII, 795 pp., Springer, New York.

345 Kertz, W. (1958), Ein neues Mass für die Feldstärke des erdmagnetischen äquatorialen  
346 Ringstroms, *Abh. Akad. Wiss. Göttingen Math. Phys.*, 2, 83 pp.

347 Karinen, A. and K. Mursula (2005), A new reconstruction of the  $D_{st}$  index for 1932–  
348 2002, *Ann. Geophys.*, 23(1), 475, 1432-0576/ag/2005-23-475.

349 Le Sager, P. and L. Svalgaard (2004), No increase of the interplanetary electric field  
350 since 1926, *J. Geophys. Res.*, 109, A07106, doi:10.1029/2004JA010411.

351 Lockwood, M., R. Stamper, and M. N. Wild (1999), A doubling of the Sun's coronal  
352 magnetic field during the past 100 years, *Nature*, 399, 437, doi:10.1038/20867.

353 Lockwood, M., A. P. Rouillard, I. Finch, and R. Stamper (2006), Comment on “The *IDV*-  
354 *index*: Its derivation and use in inferring long-term variations of the interplanetary  
355 magnetic field strength” by Leif Svalgaard and Edward W. Cliver, *J. Geophys. Res.*, 111,  
356 A09109, doi:10.1029/2006JA011640.

357 Lockwood, M., A.P. Rouillard, and I. D. Finch (2009), The Rise and Fall of Open Solar  
358 Flux during the Current Grand Solar Maximum, *Ap. J.*, 700, 937 doi:10.1088/0004-  
359 637X/700/2/937.

360 Love, J. J. (2006), Personal communication.

361 Love, J. J. (2007), A Continuous Long-Term Record of Magnetic-Storm Occurrence and  
362 Intensity, *Eos Trans. AGU*, 88(23), AGU Spring Meeting, 2007, Abstract SH54B-03.

363 Mayaud, P. N. (1980), Derivation, Meaning, and Use of Geomagnetic Indices, *Geophys.*  
364 *Monogr. Ser.*, vol. 22, 154 pp., AGU, Washington D.C.

365 McCracken, K. G. (2007), The heliomagnetic field near Earth, 1428-2005, *J. Geophys.*  
366 *Res.*, 112, A09106, doi:10.1029/2006JA012119.

367 Moos, N. A. F. (1910), *Colaba Magnetic Data, 1846 to 1905*, 2, *The Phenomenon and its*  
368 *Discussion*, 782 pp., Central Govt. Press, Bombay.

369 Muscheler, R., F. Joos, J. Beer, S. A. Müller, M. Vonmoos, and I. Snowball (2007), Solar  
370 Activity during the last 1000 yr inferred from radionuclide records, *Quart. Sci. Rev.* 26,  
371 82.

372 Owens, M. J., N. U. Crooker, N. A. Schwadron, T. S. Horbury, S. Yashiro, H. Xie, O. C.  
373 St. Cyr, and N. Gopalswamy (2008), Conservation of open solar magnetic flux and the  
374 floor in the heliospheric magnetic field, *Geophys. Res. Lettrs.*, 35, L20108,  
375 doi:/10.1029/2008GL035813.

376 Rouillard, A. P., M. Lockwood, and I. Finch (2007), Centennial changes in the solar wind  
377 speed and in the open solar flux, *J. Geophys. Res.*, 112, A05103,  
378 doi:10.1029/2006JA012130.

379 Schatten, K. H., P. H. Scherrer, L. Svalgaard, and J. M. Wilcox (1978), Using Dynamo  
380 Theory to Predict the Sunspot Number During Solar Cycle 21, *Geophys. Res. Lettrs.*,  
381 5(5), 411, doi: 10.1029/GL005i005p00411.

382 Schatten, K. H. (2005), Fair space weather for solar cycle 24, *Geophys. Res. Lettrs.*,  
383 32(21), L21106.

384 Schmidt, A. (1926), *Archiv des Erdmagnetismus*, Heft 4, Potsdam, 4.

385 Steinhilber, F., J. Beer, and C. Fröhlich (2009), Total solar irradiance during the  
386 Holocene, *Geophys. Res. Lettrs.*, 36 (19), L19704, doi:10.1029/2009GL040142.

387 Steinhilber, F., J. A. Abreu, J. Beer, and K. G. McCracken (2010), Interplanetary  
388 magnetic field during the past 9300 years inferred from cosmogenic radionuclides, *J.*  
389 *Geophys. Res.*, *115*, A01104, doi:10.1029/2009JA014193.

390 Svalgaard, L. (1977), Geomagnetic activity: dependence on solar wind parameters, in  
391 *Skylab Workshop on Coronal Holes and High Speed Streams*, ed. J. B. Zirker, Colorado  
392 Assoc. Univ. Press, Chapter 9, 371.

393 Svalgaard, L. (2005), Rederivation of Dst Index, American Geophysical Union, Fall  
394 2005, #SA12A-04, [http://www.leif.org/research/AGU%20Fall%202005%20SA12A-](http://www.leif.org/research/AGU%20Fall%202005%20SA12A-04.pdf)  
395 [04.pdf](http://www.leif.org/research/AGU%20Fall%202005%20SA12A-04.pdf)

396 Svalgaard, L. (2009), Observatory Data: a 170-year Sun-Earth Connection, in  
397 *Proceedings of the XIIIth IAGA Workshop on geomagnetic observatory instruments, data*  
398 *acquisition, and processing*: U.S. Geological Survey Open-File Report 2009-1226, ed. J.  
399 J. Love., p. 246.

400 Svalgaard, L. and E. W. Cliver (2005), The *IDV-index*: Its derivation and use in inferring  
401 long-term variations of the interplanetary magnetic field strength, *J. Geophys. Res.*, *110*,  
402 A12103, doi:10.1029/2005JA011203.

403 Svalgaard, L. and E. W. Cliver (2006), Reply to the comment by M. Lockwood et al. on  
404 “The *IDV-index*: Its derivation and use in inferring long-term variations of the  
405 interplanetary magnetic field strength”, *J. Geophys. Res.*, *111*, A09110,  
406 doi:10.1029/2006JA011678.

407 Svalgaard, L. and E. W. Cliver (2007a), A Floor in the Solar Wind Magnetic Field,  
408 *Astrophys. J.*, *661*, L203.

409 Svalgaard, L. and E. W. Cliver (2007b), Interhourly-variability index of geomagnetic  
410 activity and its use in deriving the long-term variation of solar wind speed, *J. Geophys.*  
411 *Res.*, *112*(10), A10111, doi:10.1029/2007JA012437.

412 Svalgaard, L., E. W. Cliver, and P. Le Sager (2003), Determination of interplanetary  
413 magnetic field strength, solar wind speed, and EUV irradiance, 1890–2003, in  
414 *Proceedings of ISCS 2003 Symposium: Solar Variability as an Input to the Earth's*  
415 *Environment*, Euro. Space Agency Spec. Publ., ESA SP-535, 15.

416 Wang, Y.-M. and N. R. Sheeley Jr. (2003), On the fluctuating component of the Sun's  
417 large-scale magnetic field, *Astrophys. J.*, *590*, 1111, doi:10.1086/375026.

418 Wang, Y.-M., J. L. Lean, and N. R. Sheeley Jr. (2005), Modeling the Sun's magnetic  
419 field and irradiance since 1713, *Astrophys. J.*, *625*, 522, doi:10.1086/429689.

420 Wolf, R. (1884), *Astr. Mitt. Zürich* *LXI*, 7, 1.

421 **Captions**

422 Figure 1. Adopted divisors (blue circles) to normalize *IDV* to the NGK-scale as a  
423 function of average corrected geomagnetic latitude for each station over the time of  
424 operation. For each station, different color coded symbols show what the divisor would  
425 have been for that station for years 1800 (pink pluses), 1900 (orange triangles), and 2000  
426 (red diamonds).

427 Figure 2. *IDV* calculated without any normalization or adjustments for the long-running  
428 German series (Potsdam POT–Seddin SED–Niemegk NGK; reddish curves) and the  
429 long-running French series (Parc Saint-Maur PSM–Val Joyeux VLJ–Chambon-la-Forêt  
430 CLF; greenish curves). Vertical lines show when the replacement stations went into  
431 operation and the ovals show when the yearbook values changed from being  
432 instantaneous hourly spot values to hourly means. The blue (spot values) and red (hourly  
433 means) crosses show the ratio between raw *IDVs* for KAK-Kakioka and SED-NGK.  
434 KAK changed from recording spot values to recording hourly means in 1955. There are  
435 no clear indications of changes in *IDV* due to the change in recording/reporting practice.

436 Figure 3. Raw *IDV* for the average of stations VLJ and DBN [both reporting  
437 instantaneous ‘spot’ values every hour for the 12-year intervals before 1915 and after  
438 1915] and for the average of CLH and HON [reporting spot values before 1915 (pink)  
439 and hourly mean value thereafter (blue)].

440 Figure 4 Comparison of observed *IDV* (red open squares) and synthetic *IDV* calculated  
441 from *s* using the regression equation shown (filled blue circles) derived from the

442 correlation between *IDV* and the Summed Ranges,  $s$ , of  $H$  from Colaba [Moos, 1910;  
443 page 294, table 261] for 1872-1905.

444 Figure 5. *IDV* plotted against the amplitude of the daily range,  $rY$ , of the East component  
445 [Table E2] of the geomagnetic field derived from PSM-VLJ-CLF and POT-SED-NGK,  
446 covering the interval 1883-2008 [dark blue diamonds] for which we have data for these  
447 stations. Since 1957, the number of stations contributing to *IDV* is high [ $\sim 50$ ] for every  
448 year, so the data is good. The open red squares show the same relationship for 1957-  
449 2008.

450 Figure 6. Raw *IDV* derived using night-time hourly data (blue) and daytime hourly data  
451 (red) for the French stations PSM-VLJ-CLF and the German stations POT-SED-NGK.  
452 The daytime values are  $\sim 30\%$  larger than the night-time value because of day-to-day  
453 [non-solar] variations of the regular solar variations,  $S_R$ , and have been scaled to the same  
454 average as the night-time values for easier comparison.

455 Figure 7. Yearly *IDV*-indices derived for individual stations (as given in Table 1) shown  
456 as grey curves. The red curve is a composite index calculated as the mean of the median  
457 and average values of the individual station values. This procedure may be justified by  
458 the very small difference between medians and averages (0.16 nT on average, see Figure  
459 8). The number,  $N$ , of contributing stations is shown by the thin blue curve and the  
460 corresponding number for *IDV05* as a dashed light blue line. The  $u$ -measure is  
461 considered a single station. A few station values differing more than five standard  
462 deviations from the average for a given year were omitted in calculating the average for  
463 that year. The dashed line shows *IDV* derived from the  $u$ -measure.



464 Figure 8. Average yearly values of  $IDV_{09}$  (dark blue curve) compared with median  
465 yearly values (light blue curve) and compared with published  $IDV_{05}$  (red curve).

466 Figure 9. Yearly average values of  $IDV$  and of  $D_{st}$  when it was less than zero (based on  
467  $D_{st}$  from Kyoto WDC and on  $D_{st}$  from *Love* [2006] scaled to Kyoto levels). The ‘spike’ in  
468 1909 is due to the extremely strong storm on 25 September 1909 causing loss of data at  
469 all but one station (API), giving that one data point undue influence. To guard against the  
470 influence of such sporadic extreme values, the daily values of  $IDV$  were capped at 75 nT.

471 Figure 10. Yearly average values of the HMF  $B$  inferred from the  $IDV$ -index (dark blue  
472 curve) and from the early  $u$ -measure (light blue curve) compared with in situ  
473 measurements (red curve). There is a hint of the  $\sim 100$  year Gleissberg cycle.

474 Figure 11. Comparison of HMF  $B$  determined from  $IDV$  [light blue curve using eq.(3)  
475 and dark blue curve using eq.(2)], by *Lockwood et al.* [2009, green curve], and observed  
476 by spacecraft [red curve].

477 Figure 12. Comparison between HMF  $B$  derived by *Svalgaard and Cliver* [2005] (light  
478 blue curve and open circles), this paper (dark blue curve and open circles) and HMF  $B$   
479 derived by *Lockwood et al.* [1999] (orange curve and plus-symbols), *Rouillard et al.*  
480 [2007; their point for 1901 was in error, A. Rouillard, Personal comm. 2007] (pink curve  
481 and plus symbols), and *Lockwood et al.* [2009] (red curve and plus-symbols), matched to  
482 *in situ* observations of  $B$  (black dots).

483 Figure 13. Yearly averages of near-Earth HMF  $B$  inferred by *Svalgaard and Cliver* [this  
484 paper] (blue curve  $B_{S\&C}$ ), by *Lockwood et al.* [2009] (green curve  $B_{LR\&F}$ ), observed by


485 spacecraft (red curve  $B_{\text{OBS}}$ ) compared to  $B$  inferred by *McCracken* [2007] (purple curve  
486  $B_{\text{McC}}$ ). The large arrow marks the beginning of the neutron monitor-based part of the  
487 record. One might speculate that the extremely low values during 1883-1893 are caused  
488 by the explosion of Krakatoa ejecting sulfur-rich aerosols into the stratosphere  
489 influencing the deposition of  $^{10}\text{Be}$ .


490 Figure 14. Comparison of our reconstruction of HMF  $B$  (red curve, 25 year running  
491 means) with other reconstructions as reported by *Steinhilber et al.* [2010] in their Figure  
492 7 [adapted and reproduced here], *e.g.* with their 25-year running mean of their PCA-  
493 based reconstruction [purple dashed line] of  $B$ . The oval outlines an area of disagreement  
494 for which sufficient geomagnetic data exists that may be used to resolve the discrepancy.

495 Figure 15.  $IDV$  [blue curves] and inferred HMF  $B$  [red curve; dashed line: 25-year  
496 running mean] 1880-1920 for all stations [as noted by their IAGA designations –  $10u$   
497 shown as a dashed gray line] where good geomagnetic data are available so far.


498 Table 1. Stations used for  $IDV09$ , including replacement stations due to relocation of  
499 original stations. The Corrected Geomagnetic Latitude for the year 2000 is given for  
500 illustration, but the centroid of the latitudes for the time of operation was used to estimate  
501 the Normalization Constants. Constants in *italics* were determined by an empirical fit to  
502 time-overlapping stations. For a few observatories (marked with an asterisk) weakly non-  
503 linear relationships have been used to normalize directly to NGK. A list of IAGA  
504 designations, observatory names, and other station details can be found at  
505 [http://www.geomag.bgs.ac.uk/gifs/annual\\_means.shtml](http://www.geomag.bgs.ac.uk/gifs/annual_means.shtml).

506







No Difference Between Spot and Mean Values Before and After 1915




Comparison of IDV and Colaba Summed Ranges





**IDV as a Function of Daily Range of East Component**
















Negative  $D_{st}$  and IDV measure the same effect



















Stations (IAGA Abbrev.)	Geodetic Latitude	Geodetic Longitude	Corrected Geomagnetic Latitude 2000	Divisor
BOX	58.0	39.0	53.9	1.10
ESK*	55.3	356.8	52.9	1.00
EKT,SVD,ARS	56.4	58.6	52.1	1.10
RSV,BFE	55.6	11.7	52.1	1.10
MOS	55.5	37.3	51.3	1.10
NVS	55.0	82.9	50.5	0.97
WLH,WNG	53.7	9.1	50.1	0.97
MNK	54.1	26.5	49.9	0.98
CLH,FRD	38.2	282.6	49.7	0.97
BOU	40.1	254.8	49.2	0.99
BAL	38.8	264.8	49.0	0.99
DBN,WIT	52.1	5.2	48.4	0.98
10u	52.4	13.1	48.3	1.00
POT,SED,NGK	52.1	12.7	48.0	1.00
ABN,HAD	51.0	355.5	47.8	0.99
BEL	51.8	20.8	47.5	1.01
IRT	52.2	104.5	47.0	1.02
TKT	41.3	69.6	46.5	1.08
PET	53.1	158.6	46.3	1.02
DOU	50.1	4.6	46.0	1.02
LVV	49.9	23.8	45.3	1.04
PSM,VLJ,CLF	48.0	2.3	43.6	1.04
FUR	48.2	11.3	43.4	1.05
HRB	47.9	18.2	43.0	1.06
THY	46.9	17.9	41.8	1.08

YSS	47.0	142.7	39.9	1.10
TUC	32.3	249.2	39.9	1.10
AAA	43.3	76.9	38.4	1.12
TFS	42.1	44.7	37.2	1.14
MMB	43.9	144.2	36.7	1.13
AQU	42.4	13.3	36.3	1.13
BJI,BMT	40.3	116.2	34.2	1.16
SFS,EBR	40.8	0.5	34.2	1.14
COI	40.2	351.6	34.1	1.15
LNP,LZH	36.1	103.9	30.1	1.20
VQS,SJG	18.4	293.9	29.2	1.20
KAK	36.2	140.2	28.9	1.20
KNZ	35.3	140.0	27.9	1.21
HTY	33.1	139.8	25.7	1.23
SSH	31.1	121.2	24.4	1.24
KNY	31.4	130.9	24.3	1.24
HON	21.3	202.0	21.7	1.26
GUI	28.3	343.6	15.7	1.29
PHU	21.0	106.0	13.7	1.30
API	13.8	188.2	12.8	1.30
ABG	18.6	72.9	11.8	1.31
KOU	5.1	307.4	10.8	1.30
MBO	14.4	343.0	3.2	1.31
ANN	11.4	79.7	3.1	1.32
TAM	22.8	5.5	3.1	1.32
HUA	-12.1	284.7	2.1	1.32
GUA	13.6	144.9	1.0	1.32
TRD	8.5	77.0	0.4	1.32

AAE	9.0	38.8	-1.3	1.32
BNG	4.4	18.6	-2.2	1.32
ASC	-7.5	345.6	-7.9	1.32
BTV	-6.2	106.8	-15.8	1.29
PPT	-17.6	210.4	-16.4	1.29
VSS	-22.4	316.4	-16.5	1.30
PIL	-31.7	296.1	-18.6	1.28
TAN	-18.9	47.6	-29.1	1.20
TSU	-19.2	17.7	-30.0	1.20
HBK	-22.9	27.7	-33.6	1.17
CTO,HER	-34.4	19.2	-42.3	1.09
WAT,GNA	-31.8	116.0	-44.4	1.05
TOO,CNB	-35.3	149.4	-45.8	1.04
TRW	-43.3	19.0	-47.8	1.02
AMS*	-37.8	77.6	-49.1	1.00
AIA	-65.2	295.7	-49.8	1.20
AML,EYR	-43.4	172.4	-50.3	0.97
CZT	-46.4	51.9	-53.1	1.10

		IDV	Obs			
Year	IDV09	HMF <i>B</i>	HMF <i>B</i>			
				1861.5	12.20	7.50
				1862.5	10.00	6.51
1835.5	11.60	7.23		1863.5	9.40	6.23
1836.5	16.30	9.30		1864.5	8.40	5.76
1837.5	16.00	9.17		1865.5	7.90	5.53
1838.5	16.80	9.51		1866.5	7.30	5.24
1839.5	14.00	8.30		1867.5	6.80	5.00
1840.5	12.20	7.50		1868.5	8.10	5.62
1841.5	10.10	6.55		1869.5	11.10	7.01
1842.5	9.00	6.05		1870.5	16.70	9.47
1843.5	8.90	6.00		1871.5	16.00	9.17
1844.5	8.50	5.81		1872.5	14.60	8.56
1845.5	9.50	6.28		1873.5	9.90	6.46
1846.5	10.10	6.55		1874.5	9.50	6.28
1847.5	11.40	7.14		1875.5	7.20	5.19
1848.5	13.10	7.90		1876.5	6.00	4.61
1849.5	12.00	7.41		1877.5	6.60	4.90
1850.5	9.90	6.46		1878.5	5.80	4.51
1851.5	9.00	6.05		1879.5	6.00	4.61
1852.5	7.60	5.39		1880.5	7.89	5.53
1853.5	7.80	5.48		1881.5	8.69	5.90
1854.5	7.60	5.39		1882.5	12.47	7.62
1855.5	5.80	4.51		1883.5	10.20	6.60
1856.5	6.60	4.90		1884.5	9.36	6.22
1857.5	7.20	5.19		1885.5	9.57	6.22
1858.5	10.60	6.78		1886.5	9.14	6.11
1859.5	14.30	8.43		1887.5	7.75	5.46
1860.5	13.50	8.08		1888.5	7.27	5.23

1889.5	6.99	5.09	1917.5	10.95	6.94
1890.5	6.22	4.72	1918.5	10.97	6.95
1891.5	8.60	5.86	1919.5	11.57	7.22
1892.5	14.02	8.31	1920.5	10.28	6.64
1893.5	10.79	6.87	1921.5	8.97	6.03
1894.5	13.12	7.91	1922.5	7.74	5.45
1895.5	9.95	6.48	1923.5	6.17	4.70
1896.5	10.48	6.73	1924.5	6.89	5.04
1897.5	8.71	5.91	1925.5	8.05	5.60
1898.5	8.98	6.04	1926.5	10.69	6.82
1899.5	7.06	5.13	1927.5	9.29	6.18
1900.5	5.75	4.49	1928.5	9.69	6.37
1901.5	4.90	4.06	1929.5	9.64	6.34
1902.5	5.04	4.13	1930.5	10.22	6.61
1903.5	7.03	5.11	1931.5	7.38	5.28
1904.5	7.54	5.36	1932.5	7.22	5.21
1905.5	8.62	5.87	1933.5	6.96	5.08
1906.5	7.49	5.33	1934.5	6.83	5.02
1907.5	8.83	5.97	1935.5	7.75	5.46
1908.5	9.45	6.26	1936.5	8.81	5.96
1909.5	9.95	6.48	1937.5	12.11	7.46
1910.5	8.10	5.63	1938.5	14.02	8.31
1911.5	7.08	5.14	1939.5	12.79	7.77
1912.5	5.69	4.46	1940.5	12.48	7.63
1913.5	5.15	4.18	1941.5	12.10	7.46
1914.5	6.22	4.72	1942.5	9.57	6.31
1915.5	8.09	5.62	1943.5	8.97	6.03
1916.5	9.19	6.13	1944.5	8.28	5.71

1945.5	8.75	5.93		1973.5	9.28	6.18	6.35
1946.5	14.33	8.44		1974.5	9.18	6.13	6.63
1947.5	13.85	8.24		1975.5	8.15	5.65	5.82
1948.5	10.87	6.91		1976.5	8.70	5.91	5.45
1949.5	13.55	8.10		1977.5	8.96	6.03	5.85
1950.5	12.56	7.66		1978.5	12.32	7.56	7.08
1951.5	12.46	7.62		1979.5	11.78	7.32	7.59
1952.5	10.97	6.95		1980.5	10.51	6.74	6.98
1953.5	8.90	6.00		1981.5	13.78	8.20	7.84
1954.5	7.46	5.32		1982.5	15.25	8.84	8.81
1955.5	8.69	5.90		1983.5	11.60	7.23	7.61
1956.5	13.38	8.02		1984.5	10.50	6.74	7.32
1957.5	16.65	9.45		1985.5	9.06	6.07	5.89
1958.5	15.42	8.92		1986.5	8.80	5.95	5.74
1959.5	14.39	8.47		1987.5	8.20	5.67	6.09
1960.5	15.87	9.11		1988.5	10.21	6.61	7.30
1961.5	11.49	7.18		1989.5	16.74	9.48	8.15
1962.5	8.62	5.87		1990.5	12.84	7.79	7.29
1963.5	8.06	5.60	5.45	1991.5	15.77	9.07	9.34
1964.5	7.19	5.19	5.12	1992.5	12.87	7.80	8.25
1965.5	6.93	5.07	5.06	1993.5	10.08	6.54	6.59
1966.5	7.88	5.52	6.00	1994.5	9.06	6.07	6.15
1967.5	10.30	6.65	6.36	1995.5	9.08	6.08	5.72
1968.5	9.47	6.26	6.19	1996.5	6.76	4.98	5.11
1969.5	9.39	6.23	6.05	1997.5	8.06	5.60	5.51
1970.5	10.12	6.56	6.35	1998.5	10.34	6.66	6.89
1971.5	8.84	5.97	6.00	1999.5	9.82	6.42	6.91
1972.5	9.49	6.27	6.38	2000.5	13.36	8.02	7.18

2001.5	13.44	8.05	6.94	2006.5	7.22	5.21	5.03
2002.5	10.90	6.92	7.64	2007.5	5.96	4.59	4.48
2003.5	12.51	7.64	7.60	2008.5	5.29	4.25	4.23
2004.5	9.42	6.24	6.53	2009.5	5.04	4.13	4.05
2005.5	9.44	6.25	6.25	2010.2	5.50	4.45	4.95