

The Effect of Weighting in Counting Sunspots

Leif Svalgaard

HEPL, Stanford University

SSN-Workshop, Sunspot, NM, Sept. 2011

Waldmeier's Description of What he Believed was the Counting Method

CHANGES TO THE COUNTING METHOD

Since Rudolph Wolf began the sunspot measurement, he set the standard. And although he counted each spot regardless of its size, he failed to include those smallest spots visible only under a stable atmosphere. Around 1882 Wolf's successors permanently changed the counting method in two ways to compensate for the large variation in spot size:

- (1) by including the smallest spots visible under an atmosphere of constant transparency and
- (2) by weighting spots with penumbrae according to their size and umbral structure.

Waldmeier, 1961

I believe (2) is **incorrect**, having read *all* Wolf's, Wolfer's, and Brunner's papers and not found any such description. Waldmeier may have believed that the spots were weighted by size and carried that belief into his count. And (1) is clearly not practical because seeing varies.

Waldmeier's Own Description of his [?] Counting Method

Astronomische Mitteilungen der Eidgenössischen Sternwarte Zürich
Nr. 285

1968

Die Beziehung zwischen der Sonnenfleck-
relativzahl und der Gruppenzahl

Von

M. WALDMEIER

Hofflecken handelte. Später wurden den Flecken entsprechend ihrer Größe Gewichte erteilt: Ein punktförmiger Fleck wird einfach gezählt, ein größerer, jedoch nicht mit Penumbra versehener Fleck erhält das statistische Gewicht 2, ein kleiner Hoffleck 3, ein größerer 5. Die Gruppen- und

“A spot like a fine point is counted as one spot; a larger spot, but still without penumbra, gets the statistical weight 2, a smallish spot with penumbra gets 3, and a larger one gets 5.” Presumably there would be spots with weight 4, too.

After Waldmeier took over the production of the sunspot series he stated { *100 Jahre Sonnenfleckenstatistik, Astron. Mitt. Eid. Sternw. Zürich, 152, 1948* }:

[...] Allerdings hat Wolfer, während seiner Assistentenzeit 1877-1893 eine andere Zählweise verwendet [...] dass die Hofflecken, die bei Wolf nur als **ein** Fleck galten, je nach ihrer Grösse und Unterteilung mehrfach gezählt werden.

([...] “Though Wolfer used an different counting method during his tenure as assistant 1877-1893 [...] that spots with penumbra, that by Wolf was counted as **one** spot, would be counted multiple times according to size and complexity”).

Kopecký et al. (*On the relative inhomogeneity of long-term series of sunspot indices, Bull. Astron. Inst. Czech., 31, 267-283, 1980*) essentially quote Waldmeier with a twist:

[...] beginning with Wolfer, a “modified” method of calculating the number of sunspots, but **without mentioning it** {matches that I have not seen any either}, is being used in Zürich.

They speculate that perhaps using the new Zürich classification of groups might introduce an inhomogeneity, quoting Zelenka [Memorandum and personal communication].

This 'modified' counting method is still in use at the reference station Locarno used by SIDC . As a typical example we take the drawing made at Locarno on 21st October, 2010 [next slide]. Three sunspot groups are visible, numbered by Locarno as 102, 104, and 107, corresponding to NOAA active region numbers 11113, 11115, and 11117.

From Hathaway's list we get the areas of those spots:

Year	M	D.	UT	NOAA	Loc#	Area (obs.)
2010	10	21.	50	11113	102	134 μH
2010	10	21.	50	11115	104	223 μH
2010	10	21.	50	11117	107	104 μH

No. 245

2010.X.21.333

08:00 T.U.

Osservatore: M. CAGNONI

Immagini: 3-4 (SIDC: 2-3)

$\Delta p = -25'$

SPECOLA SOLARE TICINESE
LOCARNO MONTI

$L_o = 119.9$

$B_o = +5.4$

$P_o = +25.8$

g	f	t	B	L	Δ
102	5	J	+16'		
104	3	J	-25'		
107	3	J	+23'		
3	11				

g	f	t	B	L
102	5	J	+16'	
104	3	J	-25'	
107	3	J	+23'	
3	11			

Drawing from Locarno 21 October, 2010 showing the three Locarno Regions 102, 104, and 107. The table gives the weight assigned to each group.


An insert (red border) shows the regions as observed at MWO on the 17th October (no observation the 21st).

The raw sunspot number reported by Locarno (upper right-hand table) was $3 \times 10 + 11 = 41$, which with Locarno's standard k-factor of 0.60 translates to a reduced relative sunspot number on the Wolf scale of $0.6 \times 41 = 25$ which is indeed what SIDC reported for that day.

Wolf would have reported $3 \times 10 + 3 = 33$, so rough indication of the effect of weighting would be $41/33 = 1.24$

From Hathaway's list we get the areas of those spots:

Year	M	D.	UT	NOAA	Loc#	Area (obs.)
2010	10	21.	50	11113	102	134 μH
2010	10	21.	50	11115	104	223 μH
2010	10	21.	50	11117	107	104 μH



- Note this spot of the same size: -
 1920 11 21.55 9263 *MWO* 223 μH

Up until Waldmeier [who discontinued this!] the Zürich observers recorded their raw data for each day in this format

“Group Count•Total Spot Count”

Sonnenfleckenbeobachtungen im Jahre 1849.


	I.	II.	III.	IV.	V.	VI.	VII.	VIII.	IX.	X.	XI.	XII.
1	9.31	3. 6	4. -	10.70	9.30	8.48	4.13	4 15	7.64	8.10	5.16	—
2	9.34	7.40	5. -	7. -	9.40	9.64	3. 3	6.18	5.35	7.10	7.41	8. 9
3	15. -	2. -	6.12	10.38	5.12	8.50	3. 6	6.15	4.27	3. 4	3.10	8.17
4	9.31	7.27	7.15	12.58	7.45	10.50	3 10	4.12	5.41	2. 3	4.31	—
5	9. -	9.22	2. -	8.20	8 50	8.45	7. -	5.20	1. 1	1. 2	—	9.47
6	8. -	10 34	7.24	10.60	7.38	7.45	4. 8	4.18	6.25	4. 6	—	2. 2
7	—	3. -	3. -	8.24	1. -	5. -	5.10	3.20	7.48	—	6.22	—
8	8.28	10.21	4. -	6.20	6.20	5.12	6.15	3.15	5.38	5.16	7.35	—
9	8.30	10.35	3. -	9.45	6.25	3. -	7.20	4.14	7.50	5.26	6.20	—

To calculate the relative sunspot number, e.g. on April 4th, one performs

$$R = k * (10 * 12 + 58) = 178$$


where the scale factor k is 1.00 for Wolf himself.

So, now back to the MWO spot on 21st Nov. 1920 that had the same size as Locarno 104 [which was counted as three spots or 1 spot with weight of 3.]



The insert shows a similar group observed at MWO on 5th Nov., 1922. For both groups, Wolfer should have recorded the observation as 1.3 if he had used the weighting scheme, but they were recorded as 1.1, clearly counting the large spots only once (*thus with no weighting*). The recorded (in the historical record) Zürich sunspot number was 7 $\{=0.6 \times (10+1)\}$ on both those days, consistent with no weighting.

Other Observatory Drawings Show Similar Results, e.g. Haynald (Kalocsa, Hungary):



This spot should have been counted with weight 3, so the recorded value should have been 1.3, if Wolfer had applied the weighting, which he obviously didn't

There are many other such examples, (e.g. 16th September, 1922 and 3rd March, 1924 for which MWO drawings are readily available). **We thus consider it established that Wolfer (and by extension [?] the other observers before Waldmeier) did not apply the weighting scheme contrary to Waldmeier's assertion.**

This is consistent with the fact that nowhere in Wolf's and Wolfer's otherwise meticulous yearly reports in the *Mittheilungen über Sonnenflecken* series is there any mention of a weighting scheme. Furthermore, Wolf was still very much alive in 1882 and in charge of things, and was not 'succeeded' at that time. Waldmeier himself was an assistant to Brunner in 1936 and performed routine daily observations with the rest of the team so should have known what the rules were. There is a mystery lurking here. Perhaps the Archives [in Zürich? Or the microfilm in Brussels] will provide a resolution of this conundrum.

What Do the Observers at Locarno Say About the Weighting Scheme:



Sergio Cortesi started in 1957, still at it, and in a sense is the *real* keeper of the SSN, as SIDC normalizes everybody's count to match Sergio's

“For sure the main goal of the former directors of the observatory in Zürich was to maintain the coherence and stability of the Wolf number, and changes in the method were not done just as fun. I can figure out that they gave a lot of importance to verify their method of counting. Nevertheless the decision to maintain as “**secret**” the true way to count is for sure source of problems now!”

(email 6-22-2011 from Michele Bianda, IRSOL, Locarno)

Estimating Unweighted Sunspot Count From Locarno Drawings

I look at the drawing of a group and from experience [I have looked at thousands of spots, 42025 at last count] on Locarno's drawings going back many years], assign a weight to each spot, then subtract the weight from the count given for the group and add 1 for the spot.

Example 1: A group has four spots on the drawing, one is large with weight 3, one is medium with weight 2 and two are small with weight 1. The total count given by Locarno was 6. That tells me that one of the small spots was not counted [otherwise the total would have been $3+2+1+1 = 7$]. So, I subtract 3, 2, and 1 from their total: $6 - 3 - 2 - 1 = 0$ and add 1 for each spot for a total of 3 as the unweighted count.

Example 2: Most of the time it is enough just to count the spots:

136	3	J	+10	
138	2	A	+17	
139	5	C	-8	2
4	65			


2004-8-12

3 2
139

More Examples


Just counting the spots regardless of size.
How does Marco get 20 for group 162?

Often there are more spots on the drawing than were included in the visual count at the eyepiece:



No. 187
 2011.VIII.16.281
 06:45 T.U.
 Osservatore: M. CAGNOTI
 Immagini: 2 (SIDC: 4)

g	f	t	B	L	Δ
162	20	0	+15'		7
163	3	8	+18'		1
164	3	8	-21'		1
<hr/>					
3	26				9
<hr/>					



2006.IV.20.313
 7.30 T.U.
 Osservatore: S. Cortesi
 Immagini: 2-3

g	f	t	B	L	
30	4	B	-1		
<hr/>					
1	4				
<hr/>					

3

Difficult Cases



2005. IX. 15. 333
 8.00 T.U.
 Osservatore: S. Cortesi
 Immagini: 3

g	f	t	B	L	
105	58	E	-11		36
1	58				

105

74 spots

3,2,3,2,2,3,2,3,3,3,3,3: sum 35, 58-35+13 spots = 36



g	f	t	B	L	△
134	55	E	-13		40
136	3	J	+10		1
138	2	A	+17		1
139	5	C	-8		2
4	65				44

2004-08-12 (group 134)

Examples of Spots Not Counted


2003.III.11.385

9.15 T.U.

Osservatore: S. Cortesi

Immagini: 2 schiarite

$\Delta p = +23.7$



70

g	f	t	B	L
57	4	H	+12	1
65	4	B	-13	4
66	12	G	+6	
68	3	B	-24	
69	13	D	-12	
70	6	B	+15	6
6	42			

Details of My Analysis

(covering 2003-2011 so far)

				G	S	G	S	R	R			R	
				Loc	Loc	Leif	Leif	Loc	Leif	Loc/Leif	Obs	SIDC	<i>k</i> Loc
2011	8	28	2011.657	5	22	5	15	72	65	1.1077	am	49	0.6806
2011	8	29	2011.660	6	19	6	12	79	72	1.0972	mc	43	0.5443
2011	8	30	2011.662	7	22	7	17	92	87	1.0575	mc	69	0.7500
2011	8	31	2011.665	9	57	9	35	147	125	1.1760	mc	96	0.6531
2011	9	1	2011.666	7	59	7	33	129	103	1.2524	mb	85	0.6589
2011	9	2	2011.669	8	72	8	52	152	132	1.1515	mc	91	0.5987
2011	9	3	2011.672	8	74	8	49	154	129	1.1938	mc	95	0.6169
2011	9	4	2011.674									70	
2011	9	5	2011.677	6	70	6	53	130	113	1.1504	mc	76	0.5846
2011	9	6	2011.680	5	37	5	25	87	75	1.1600	mc	58	0.6667
2011	9	7	2011.683									49	
2011	9	8	2011.685	3	24	3	12	54	42	1.2857	mc	36	0.6667
2011	9	9	2011.688	4	38	4	23	78	63	1.2381	mc	46	0.5897
2011	9	10	2011.691	5	36	5	25	86	75	1.1467	am	52	0.6047
2011	9	11	2011.694	7	44	7	28	114	98	1.1633	am	59	0.5175
										1.1677			0.6303

The SIDC numbers are preliminary and are updated as needed

Double-Blind Test

Email from Leif Svalgaard


Sat, Jun 18, 2011 at 9:26 PM

Dear Everybody,

As you may know we are holding a sunspot workshop at Sunspot, New Mexico in September. For this I would like to propose a simple test, that hopefully should not put a great extra burden on everybody. I ask that the observer for each day writes down somewhere what the actual number of spots counted was without the weighting, but without telling me. Then in September you let me know what the counts for [rest of] June, July, and August were. This allows me to calibrate my method of guessing what your count was. It is, of course, important that the test be blind, that I do not know until September what you all are counting. I hope this will be possible.

My modest proposal was met with fierce resistance from everybody [incl. Frédéric], but since I persisted in being a pest, I finally got Locarno to go along

Current Status of the Test



2nd degree fit


S	Sw	Sw/S
10	14.74	1.4737
25	34.83	1.3933
50	64.81	1.2961
75	90.38	1.2051
100	111.55	1.1155

For typical number of spots the weighting increases the 'count' of the spots by 30-50%

For the limited data for August 2011 Marco Cagnotti and Leif Svalgaard agree quite well with no significant difference. The test should continue as activity increases in the coming months.



Comparison of 'Relative Numbers'




But we are interested in the effect on the SSN where the group count will dilute the effect by about a factor of two.

For Aug. 2011 the result is at right. There is no real difference between Marco and Leif.

We take this a [preliminary] justification for my determination of the influence of weighting on the Locarno [and by extension on the Zürich and International] sunspot numbers



Weight Factor and Sunspot Number

(The weight factor is how much the weighting increases the sunspot number)



Daily values

Temporal Evolution of Weight Factor



Daily values

The Average Weight Factor

$$1.13 + 0.00040 * R$$


0.6176Avg

R=100

1.17

0.6088Med


	slope					inv. Slope			
all	0.8722	<Rloc>	<Rleif>	ratio	<Rsidc>	1.1465	count	Loc	k loc
2011.4	0.8691	70.29	61.36	0.8728	42.84	1.1506	211	0.6094	
2010.5	0.8767	28.30	24.96	0.8822	16.47	1.1406	285	0.5819	
2009.5	0.8945	4.74	4.32	0.9119	3.12	1.1179	309	0.6570	
2008.5	0.8807	4.00	3.64	0.9107	2.85	1.1355	297	0.7137	
2007.5	0.8801	12.33	10.90	0.8842	7.50	1.1362	332	0.6088	
2006.5	0.8814	24.55	21.89	0.8919	15.22	1.1346	312	0.6200	
2005.5	0.8662	50.37	43.80	0.8696	29.83	1.1545	318	0.5922	
2004.5	0.8838	68.63	60.50	0.8816	40.45	1.1315	303	0.5894	
2003.5	0.8654	108.69	93.83	0.8632	63.71	1.1555	190	0.5861	



For yearly values there is an approximately (but weak) linear relation between the weight factor and the sunspot number. For a typical R of 100, the weighting increases the sunspot number by 17%.

How Many Groups?


(The Waldmeier Classification May lead to Better [larger] Determination of Groups)



Counting Groups

- This deserves a full study. I have only done some preliminary work on this, but estimate that the effect amounts to a few percent only, perhaps 3% [?]
- This would increase the ‘Waldmeier Jump’ to about 20%
- My suggested solution is to increase all pre-Waldmeier SSNs by 20%, rather than decrease the modern counts which may be used in operational programs


Can we see the Effect in the Data?



We can compute the ratio R_z/R_g [staying away from small values] for some decades on either side of the start of Waldmeier's tenure, assuming that R_g derived from the RGO data has no trend over that interval.

There is a clear discontinuity corresponding to a jump of a factor of 1.18 between 1945 and 1946. This compares favorably with the estimated size of the increase due to the weighting [with perhaps a very small additional influence from a greater group count]

Sunspot Areas vs. Rz




The relationship between sunspot number and sunspot area [SA, Balmaceda] is not linear, but can be made linear raising SA to the power of 0.732. Then taking the ratio makes sense.

Pink squares show the ratios for SA exceeding 1000 micro-hemispheres

Clear change in the relationship around 1945

Quantifying the Waldmeier 'Jump'

Histogram Ratios



1874-1944 0.3244


1945-2000 0.3921

Waldmeier Jump

$0.3921/0.3244 = 1.212$


Plotting Histograms of the ratio $Rz/SA^{0.732}$

Illustrating that Observed Rz after 1945 is Higher than Deduced from Sunspot Areas






Ca II K-line Data Scaled to Rz shows similar Jump in Rz Sunspot Number after 1945

From ~40,000 CaK spectroheliograms from the 60-foot tower at Mount Wilson between 1915 and 1985, a daily index of the fractional area of the visible solar disk occupied by plages and active network has been constructed [Bertello et al., 2008]. Monthly averages of this index is strongly correlated with the sunspot number $SSN = 27235 \text{ CaK} - 67.14$ [before 1945].




Waldmeier's Sunspot Number 19% higher than Brunner's from Ca II K-line



Digression: Change of MWO MPSI Calibration in 1982 after Upgrade of Instrument



The Amplitude of the Diurnal Variation [from many stations] shows the same Change ~1945



foF2



FIGS. 1 AND 2—PLOT OF 12-MONTH RUNNING AVERAGE OF MONTHLY MEDIAN $f^{\circ}F_2$ AGAINST 12-MONTH RUNNING AVERAGE OF MONTHLY ZURICH SUNSPOT NUMBER, LOCAL TIME

F2-layer critical frequency. This is the maximum radio frequency that can be reflected by the F2-region of the ionosphere at vertical incidence (that is, when the signal is transmitted straight up into the ionosphere). And has been found to have a profound solar cycle dependence.

The shift in SSN to bring the curves to overlap is 21%

So, many lines of evidence point to an about 20% Waldmeier Weighting Effect